APS/123-QED

arXiv:quant-ph/0502073v3 14 Jul 2005

Dwell times for transmission and reflection

N. L. Chuprikov*
Tomsk State Pedagogical University, 634041, Tomsk, Russia†

(Dated: February 1, 2008)

As was shown in quant-ph/0405028, the state of a tunneling particle can be uniquely presented as a coherent superposition of two states to describe alternative sub-processes, transmission and reflection. In this paper, on the basis of the stationary wave functions for these sub-processes, we give new definitions of the dwell times for transmission and reflection. In the case of rectangular potential barriers the dwell times are obtained explicitly. In contrast with the well-known Büttiker's dwell-time, our dwell time for transmission increases exponentially, for the under-barrier tunneling, with increasing the barrier's width. By our approach the well-known Hartman effect is rather an artifact resulted from an improper interpretation of the wave-packet tunneling and experimental data, but not a real physical effect accompanying the tunneling phenomenon.

PACS numbers: 03.65.Ca, 03.65.Xp

I. INTRODUCTION

In this paper we continue our study [1] of the temporal aspects of tunneling considered in our approach as a combined stochastic process to consist from two alternative elementary ones, transmission and reflection, evolved coherently. In [1] we have introduced the transmission and reflection times provided that a particle is in a localized state. These time scales were extracted from the time dependence of the average positions of transmitted and reflected particles.

It is evident that this way of timing a particle is not applicable in the limiting case, when the particle is in some stationary state described by a non-normalized wave function. This case is of great importance, since it is connected to the scattering of uniform beams of identically prepared particles, noninteracting with each other. To find a time spent by transmitted and reflected particles in the barrier region, in the stationary scattering problem, is the main aim of the present paper.

II. STATIONARY WAVE FUNCTIONS FOR TRANSMISSION AND REFLECTION IN THE CASE OF RECTANGULAR POTENTIAL BARRIERS

How to obtain the wave functions for transmission and reflection, for a given potential and initial wave function, is presented in [1]. Here we expose both the functions in the form to be more suitable for our purpose.

Let setting the scattering problem be such as in [1]. In particular, we will suppose that the rectangular potential barrier of height V_0 is localized in the interval $[a,b]$. Then the wave functions for transmission, $\Psi_{tr}(x;E)$, and reflection, $\Psi_{ref}(x;E)$, can be written for $E\le V_0$ as follows (E is the energy of a particle).

For $x\le a$

$$\Psi_{tr}=A_{in}^{tr}e^{ikx};\quad \Psi_{ref}=A_{in}^{ref}e^{ikx}+b_{out}e^{-ik(x-2a)};\tag{1}$$

for $a\le x\le x_c$

$$\begin{aligned}\Psi_{tr}&=a_{tr}^{l}\sinh(\beta)+b_{tr}^{l}\cosh(\beta);\\ \Psi_{ref}&=a_{ref}^{l}\sinh(\beta);\end{aligned}\tag{2}$$

for $x_c\le x\le b$

$$\Psi_{tr}=a_{tr}^{r}\sinh(\beta)+b_{tr}^{r}\cosh(\beta);\quad \Psi_{ref}=0;\tag{3}$$

for $x\ge b$

$$\Psi_{tr}=a_{uot}e^{ik(x-d)};\quad \Psi_{ref}=0;\tag{4}$$

where $d=b-a$, $x_c=(b+a)/2$, $\beta=\kappa(x-x_c)$,$k=\sqrt{2mE/\hbar^2}$, $\kappa=\sqrt{2m(V_0-E)/\hbar^2}$; m is the mass of a particle. Besides,

$$A_{in}^{tr}=a_{out}^{*}\left(a_{out}+b_{out}\right),$$

$$A_{in}^{ref}=b_{out}\left(b_{out}^{*}-a_{out}^{*}\right);$$

$$a_{tr}^{l}=\frac{i}{\kappa}PA_{in}^{tr}e^{ika},\quad b_{tr}^{l}=\frac{1}{\kappa}QA_{in}^{tr}e^{ika},$$

$$a_{tr}^{r}=\frac{i}{\kappa}P^{*}a_{out}e^{ika},\quad b_{tr}^{r}=\frac{1}{\kappa}Q^{*}a_{out}e^{ika},$$

$$a_{ref}^{l}=\frac{i}{\kappa}\left(PA_{in}^{ref}-P^{*}b_{out}\right)e^{ika};$$

$$a_{out}=\frac{1}{2}\left(\frac{P}{P^{*}}+\frac{Q}{Q^{*}}\right);\quad b_{out}=\frac{1}{2}\left(\frac{P}{P^{*}}-\frac{Q}{Q^{*}}\right).$$

---

[*] Electronic address: chnl@tspu.edu.ru

[†] Also at Physics Department, Tomsk State University.

In its turn,

$$P = k\cosh(\varphi) - i\kappa\sinh(\varphi),$$

$$Q = \kappa\cosh(\varphi) + ik\sinh(\varphi); \quad \varphi = \frac{\kappa d}{2}.$$

One can easily check that

$$\frac{a_{out}}{A_{in}^{tr}} = -\frac{b_{out}}{A_{in}^{ref}} = \frac{Q}{Q^*};$$

and $|a_{out}|^2 = T$, $|b_{out}|^2 = R$ where T and R are the coefficients of transmission and reflection, respectively ($T + R = 1$); $A_{in}^{tr} + A_{in}^{ref} = 1$. Thus, $b_{tr}^l = b_{tr}^r = b_{tr}$.

III. DWELL TIME FOR TRANSMISSION

Note, both the stationary wave functions are nonnormalized and, hence, the standard timing procedure becomes usefulness in timing a particle with the well defined energy. At the same time the properties of these wave functions provide another natural way for timing a particle in the barrier region.

Indeed, the density of the probability flux, I_{tr}, for $\Psi_{tr}(x;E)$ is everywhere constant and equals to $\frac{\hbar k}{m}T$. This means that the velocity of this flux, v_{tr}, can be written as

$$v_{tr}(x) = \frac{I_{tr}}{|\Psi_{tr}(x;E)|^2}.$$

Taking into account Exps. (1) - (4) for $\Psi_{tr}(x;E)$, we see that outside the barrier region $v_{tr} = \hbar k/m$. In the barrier region, the velocity of the flux decreases exponentially when it approaches the midpoint x_c. One can easily show that $|\Psi_{tr}(a;E)| = |\Psi_{tr}(x;E)| = \sqrt{T}$, but $|\Psi_{tr}(x_c;E)| = \sqrt{T}|Q|/\kappa$.

Thus, any selected infinitesimal element of the flux passes the barrier region for the time τ_{dwell}^{tr}, where

$$\tau_{dwell}^{tr} = \frac{1}{I_{tr}}\int_a^b |\Psi_{tr}(x;E)|^2 dx. \tag{5}$$

One can easily show that

$$\int_a^b |\Psi_{tr}(x;E)|^2 dx = \frac{d}{4}\left(2|b_{tr}|^2 - |a_{tr}^r|^2 - |a_{tr}^l|^2\right) + \frac{\sinh(2\varphi)}{4\kappa}\left(2|b_{tr}|^2 + |a_{tr}^r|^2 + |a_{tr}^l|^2\right) + \frac{\sinh^2(\varphi)}{\kappa} Re[(a_{tr}^r - a_{tr}^l)b_{tr}^*]. \tag{6}$$

Then, taking into account Exps. (1) - (4)), we obtain

$$2|b_{tr}|^2 - |a_{tr}^r|^2 - |a_{tr}^l|^2 = \frac{2}{\kappa^2}(\kappa^2 - k^2)T,$$

$$2|b_{tr}|^2 + |a_{tr}^r|^2 + |a_{tr}^l|^2 = \frac{2}{\kappa^2}(\kappa^2 + k^2)T\cosh(2\varphi),$$

$$Re[(a_{tr}^r - a_{tr}^l)b_{tr}^*] = -\frac{1}{\kappa^2}(\kappa^2 + k^2)T\sinh(2\varphi).$$

Eventually, we have

$$\tau_{dwell}^{tr} = \frac{m}{2\hbar k\kappa^3}\left[\left(\kappa^2 - k^2\right)\kappa d + \kappa_0^2\sinh(\kappa d)\right]; \tag{7}$$

where $\kappa_0 = \sqrt{2m|V_0|/\hbar^2}$.

Similarly, for $E \geq V_0$

$$\tau_{dwell}^{tr} = \frac{m}{2\hbar k\kappa^3}\left[\left(\kappa^2 + k^2\right)\kappa d - \theta\kappa_0^2\sin(\kappa d)\right]; \tag{8}$$

where $\kappa = \sqrt{2m(E - V_0)/\hbar^2}$; $\theta = 1$, if $V_0 > 0$; otherwise, $\theta = -1$.

IV. DWELL TIME FOR REFLECTION

As is seen, the above time scale for transmission (5) (though the way of introducing is different) coincides by form with the well-known Büttiker's dwell time (see [3]). However, unlike the latter, Exp. (5) describes only transmitted particles.

Doing in the spirit of the Büttiker's dwell-time concept, we introduce the dwell time for reflection, τ_{dwell}^{ref}, as

$$\tau_{dwell}^{ref} = \frac{1}{I_{ref}}\int_a^{x_c} |\Psi_{ref}(x)|^2 dx; \tag{9}$$

where $I_{ref} = \frac{\hbar k}{m}R$ is the incident probability flux (note, the probability flux for $\Psi_{ref}(x)$ itself is zero).

Considering the expressions for $\Psi_{ref}(x)$ one can easily show that for $E < V_0$

$$\tau_{dwell}^{ref} = \frac{mk}{\hbar\kappa}\cdot\frac{\sinh(\kappa d) - \kappa d}{\kappa^2 + \kappa_0^2\sinh^2(\kappa d/2)}; \tag{10}$$

for $E \geq V_0$

$$\tau_{dwell}^{ref} = \frac{mk}{\hbar\kappa}\cdot\frac{\kappa d - \sin(\kappa d)}{\kappa^2 + \theta\kappa_0^2\sin^2(\kappa d/2)}. \tag{11}$$

V. OUR AND BÜTTIKER'S CONCEPTS OF THE DWELL TIME: COMPARATIVE ANALYSIS

As was shown in [2], the phase-time and old dwell-time concepts, τ_{dwell}, do not imply the passage to the case of a free particle, that is, when $\kappa_0 d \to 0$. At the same time, our concept of the asymptotic times for transmission and reflection (see [1]) ensures this passage. One can easily check that the above dwell time for transmission implies this passage, too. Note, both the characteristic times are nonnegative quantities.

Let $\tau_{free}(k)$ be the time spent by a free particle in the interval $[a,b]$: $\tau_{free}(k)=\frac{md}{\hbar k}$. Then one can show that for barriers, in the limit $k\to 0$ ($\kappa_0 d\neq 0$), we have

$$\frac{\tau^{tr}_{dwell}}{\tau_{free}}=\frac{1}{2}\left[1+\frac{\sinh(\kappa_0 d)}{\kappa_0 d}\right]; \tag{12}$$

for wells,

$$\frac{\tau^{tr}_{dwell}}{\tau_{free}}=\frac{1}{2}\left[1+\frac{\sin(\kappa_0 d)}{\kappa_0 d}\right]. \tag{13}$$

Thus, not only for $k\neq 0$, but also for $k=0$, $\tau^{tr}_{dwell}\to\tau_{free}$ when $\kappa_0 d\to 0$. Remind (see [2]), the $\tau_{dwell}/\tau_{free}=0$ in the latter case.

Remind, the Büttiker's dwell time for $E<V_0$ read as (see [3])

$$\tau_{dwell}=\frac{mk}{\hbar\kappa}\cdot\frac{2\kappa d(\kappa^2-k^2)+\kappa_0^2\sinh(2\kappa d)}{4k^2\kappa^2+\kappa_0^4\sinh^2(\kappa d)}; \tag{14}$$

for $E\geq V_0$

$$\tau_{dwell}=\frac{mk}{\hbar\kappa}\cdot\frac{2\kappa d(\kappa^2+k^2)-\kappa_0^2\sin(2\kappa d)}{4k^2\kappa^2+\kappa_0^4\sin^2(\kappa d)}. \tag{15}$$

It is useful to compare formally a new dwell-time scale (to describe transmitted and reflected particles separately) with Büttiker's dwell time to describe all particles jointly.

Figs.1 and 2 show the E-dependence of $\tau^{tr}_{dwell}/\tau_{free}$ and τ_{dwell}/τ_{free} for $V_0=-0.1eV$, $d=30nm$ and $V_0=0.1eV$, $d=15nm$, respectively. Figs. 3 and 4 show the dependence of τ_{dwell}, τ^{tr}_{dwell} and τ_{free} on the barrier's width d for $V_0=0.1eV$, $E=0.11eV$ and $V_0=0.1eV$, $E=0.09eV$, respectively. In all cases, $m=0.067m_e$ where m_e is the mass of an electron.

As is seen, τ^{tr}_{dwell} and τ_{dwell} coincide at the points of resonance. However, this result is expected. Besides, $\tau^{tr}_{dwell}(E,d)$ is more smooth function than $\tau_{dwell}(E,d)$. Moreover, except for some neighborhoods of resonant points, $\tau^{tr}_{dwell}>\tau_{dwell}$.

Our results show that the most essential difference between these time scales takes place for $E<V_0$ (see Figs.2 and 4). In particular, Fig.4 shows that τ^{tr}_{dwell} increases exponentially when the width of an opaque potential barrier increases. At the same time $\tau_{dwell}(E,d)$ (as well as the well-known phase tunneling time and our asymptotic times for transmission and reflection) tends, in this case, to some constant value.

As regards τ^{ref}_{dwell} (see (10), (11)), in this limit

$$\tau^{ref}_{dwell}=\frac{2mk}{\hbar\kappa\kappa_0^2}.$$

Note also, $\tau^{ref}_{dwell}\to 0$, for wells ($V_0<0$), when $k\to 0$. However, in the case of resonance, when $\kappa_0 d=\pi+2n\pi$, $\tau^{ref}_{dwell}\to\infty$ at $k\to 0$.

VI. SCENARIO OF TUNNELING A PARTICLE THROUGH AN OPAQUE RECTANGULAR BARRIER

As is known, the just mentioned property of the phase and dwell tomes is excepted to lead to superluminal velocities of a particle in the barrier region (the well-known Hartman effect). However, we see that the behavior of τ^{tr}_{dwell}, in contrast to the Büttiker's dwell time, denies the existence of the Hartman effect for a particle with the well-defined energy. However, from our study it follows that this effect does not appear also for a particle whose initial state is described by the wave packet of any width. Fig.5 shows the expectation value of the particle's position as a function of t. It was calculated for the transmitted wave packet providing that $a=200nm$, $b=215nm$, $V_0=0.2eV$. At $t=0$ the (combined) state of the particle is described by the Gaussian wave packet peaked around $x=0$; its half-width $10nm$; the average energy of the particle $0.05eV$.

This figure displays explicitly the difference between the exact and asymptotic transmission times. While the former gives just the (average) time spent by a particle in the barrier region, the latter describes, in fact, the (average) lag or outstripping of a transmitted particle with respect to a free particle whose average velocity equals to the asymptotic average velocity of a scattered particle.

In the case considered (see Fig.5) the exact transmission time equals approximately to $0.155ps$, the asymptotic one is of $0.01ps$, and $\tau_{free}\approx 0.025ps$. As is seen, the dwell time for transmission and exact transmission time, both evidence that, though the asymptotic transmission time is small for this opaque barrier, transmitted particles spend much time in the barrier region.

Note, the possibility of accelerating a particle in the region of an opaque potential barrier, to superluminal velocities, is doubtful *a priori.* The point is that the average energy of a particle, $\overline{\hat{H}}_{tr}$, is constant for this static potential barrier. Thus, for the average kinetic ($\overline{\hat{K}}_{tr}$) and potential ($\overline{\hat{V}}_{tr}$) energies of a particle we have, for any t,

$$\overline{\hat{K}}_{tr}+\overline{\hat{V}}_{tr}=\overline{\hat{K}}_{0,tr},$$

where $\overline{\hat{K}}_{0,tr}$ is the average kinetic energy of transmitted particles at $t=0$.

It is evident that $\overline{\hat{V}}_{tr}>0$, if $V_0>0$. Hence, at the stage of scattering, $\overline{\hat{K}}_{tr}$ may only decrease. Of course, this fact does not at all mean that the average velocity of a particle, at the stage of scattering, cannot be larger than its asymptotic value. The point is that

$$\overline{\hat{K}}_{tr}=\frac{\bar{p}^2_{tr}}{2m}+\frac{\overline{(\delta p)^2}}{2m};$$

p is the particle's momentum. Thus, $\bar{p}_{tr}$ may increase due to diminishing the wave-packet's width in the momentum space. Our calculations show that this takes place near the barrier region (see Fig.5).

Thus, when approaching the left boundary of the barrier, the particles is accelerated, on the average. As a result, it may overcome the potential barrier. When the particle enters the barrier region, it almost stops. The region of an opaque potential barrier serves as a storage of particles. Behind the barrier the average velocity of particles may first exceed its own asymptotic value. However, far from the barrier, when the width of the wave packet, in the momentum space, backs to its initial (asymptotic) value, the same does the average velocity of particles.

Note, the more is the part of $\frac{\overline{(\delta p)^2}}{2m}$ in the average kinetic energy of a particle at $t = 0$, the larger may be the grows of the particle's velocity near the barrier region. However, one has to bear in mind that this velocity cannot essentially exceed the value, $\sqrt{2mV_0}/m$. So that, as it follows from our approach, tunneling a particle through an opaque potential barrier must not be accompanied by the Hartman effect.

VII. CONCLUSION

So, in our approach, the state of the whole quantum ensemble of tunneling particles is considered as a coherent superposition of the states of the subensembles of transmitted and reflected particles. For a given potential and initial state of a particle, there is an unique pair of solutions to the one-dimensional Schrödinger equation, which describe both the subensembles at all stages of tunneling. For each of these subensembles we have introduced three time scales. Two of them (the exact and asymptotic tunneling times) describe a completed scattering process. The third time scale (the dwell time) relates to the stationary tunneling problem. It should be stressed that the asymptotic (transmission and reflection) times differ from the well-known phase asymptotic times. The dwell times for transmission and reflection differ from the Büttiker's dwell time. Analysis of tunneling a particle through an opaque rectangular barrier, carried out in the framework of our approach, does not confirm the existence of the Hartman effect.

At the end we have to point to some shortcomings of the above time concepts. So, the concept of the exact tunneling times does not give explicit expressions for these quantities; besides, it is not suitable for timing a particle with the well-defined energy (this becomes more evident in the case of the reflection of wide (in x-space) wave packets, when the expectation value of the particle's position simply does not "enter" the barrier region). The main shortcoming of the dwell-time concept is that it is valid only for the stationary scattering process.

The our next purpose is to present the concept of the Larmor times for transmission and reflection. By our opinion, this concept is of great importance in solving the tunneling time problem.

[1] N. L. Chuprikov, Wave functions and characteristic times for transmission and reflection, quant-ph/0405028.

[2] N. L. Chuprikov, A comparative analysis of tunneling time concepts: Where do transmitted particles start from, on the average?, quant-ph/0501067.

[3] M. Buttiker, 1983 Phys. Rev. B **27** 6178

Figure captions

Fig. 1. τ_{dwell}/τ_{free} and $\tau_{dwell}^{tr}/\tau_{free}$ versus E, for $d = 30nm$ and $V_0 = -0.1eV$.

Fig. 2. τ_{dwell}/τ_{free} and $\tau_{dwell}^{tr}/\tau_{free}$ versus E, for $d = 15nm$ and $V_0 = 0.1eV$.

Fig. 3. τ_{dwell}, τ_{dwell}^{tr} and τ_{free} versus d, for $E = 0.11eV$ and $V_0 = 0.1eV$.

Fig. 4. τ_{dwell}, τ_{dwell}^{tr} and $/\tau_{free}$ versus d, for $E = 0.09eV$ and $V_0 = 0.1eV$.

Fig. 5. The t-dependence of the average position of transmitted particles; the initial (full) state vector represents the Gaussian wave packet peaked around the point $x = 0$, its half-width equals to $10nm$, the average kinetic particle's energy is $0.05eV$; $a = 200nm$, $b = 215nm$.


0,0
0,1
0,2
0,3
0,4
0,5
0,6
0,7
0,8
0,00
0,05
0,10
0,15
0,20
0,25
0,30
E (eV)
τ_{dwell}/τ_{free}
$\tau^{tr}_{dwell}/\tau_{free}$


0
2
4
6
8
10
12
14
16
18
20
22
0,00
0,05
0,10
0,15
0,20
0,25
0,30
E (eV)
τ_{dwell}/τ_{free}
$\tau^{tr}_{dwell}/\tau_{free}$



τ_{dwell}(ps)
τ^{tr}_{dwell}(ps)
τ_{free}(ps)
2,4
2,2
2,0
1,8
1,6
1,4
1,2
1,0
0,8
0,6
0,4
0,2
0,0
0
50
100
150
200
250
300
d (nm)


0,08
0,07
0,06
0,05
0,04
0,03
0,02
0,01
0,00
0
10
20
30
40
50
d (nm)
τ_{dwell}(ps)
τ^{tr}_{dwell}(ps)
τ_{free}(ps)



450
400
350
300
250
200
150
100
50
0
<x>$_{tr}$ (nm)
0,0
0,1
0,2
0,3
0,4
0,5
0,6
0,7
0,8
t (ps)